

Original article

Estimate the Antioxidant Capacity, Total Phenol Contents, Mineral Concentrations, and Total Carbohydrate of *Capparis Spinosa* L. (Kabbar), *Ceratonia Siliqua* L. (Kharuwb), and *Juniperus Phoenicea* L (Arar) Plants

Khadijah Al-Awjali¹, Salma Abdulsalam², Farag El-Mokasabi³, Zuhir Akrim⁴, Hamad Hasan^{*5}

¹Botany Department, Faculty of Science, Derna University, Libya

²Biology Department, Faculty of Education, Omar Al-Muhtar University, Libya

³Botany Department, Faculty of Science, Benghazi University, Libya

⁴Pharmacology and Toxicology department, Faculty of pharmacy, Omar Al-Muhtar University, Libya

⁵Chemistry Department, Faculty of Science, Omar Al-Mukhtar University, Libya

Corresponding email. hamad.dr@omu.edu.ly

Keywords:

Phytochemical. Carbohydrate, Antioxidant, Phenols, Minerals, Plants, Libya.

ABSTRACT

In this research, some of the phytochemical compounds were detected by using phytochemical screening in leaves and stems of three different plants collected from Al-Gabal Al-Ahder region, Libya, and the concentrations of anti-oxidant capacity, total phenol content, minerals, and total carbohydrate were measured for the selected plants of *Capparis spinosa* L. (Kabbar), *Ceratonia siliqua* L. (Kharuwb), and *Juniperus phoenicea* L (Arar). The spectrophotometer was used to estimate the contents of antioxidants, total phenols, and total carbohydrates, whereas the Flame photometer was used to determine the contents of sodium, calcium, and potassium. The outcomes of this study found that, the levels of anti-oxidant were ranged as follows: (222.72 -302.64 ppm) in leaves and (307.62 -319.03 ppm) in stems, while the contents of total phenols were ranged from 307.62 to 319.03 ppm in stems, the total phenols contents were ranged from 9.07 to 10.59 ppm and from 9.95 to 10.65 ppm in stems, respectively while the concentrations of carbohydrate were ranged from 0.039 to 0.155 ppm and from 0.070 to 0.123 ppm for leaves and stems, respectively. . whereas the sodium, potassium, calcium were ranged from 0.958 to 2.625 ppm in leaves and from 0.88 to 4.08 ppm in stems, for potassium the values were ranged from 3.56 to 77.16 ppm in leaves, but for stems the values of potassium were ranged from 42.96 to 76.96 ppm. While the calcium contents ranged from 0.08 to 0.28 ppm in leaves, and for stems ranged from 0.375 to 0.875 ppm in stems. The phytochemical investigation showed the present important quantiles of sterols, flavonoids, alkaloids, tannins, anthraquinones, saponins, and phenols.

Introduction

Libya has been considered to be abundant in therapeutic plants commonly utilised in traditional medicine, particularly Cyrenaica, which is home to the majority of these plants [1]. The Sylphium of antiquity was the most well-known medicinal plant that greatly enriched Cyrenaica in distant times. Due to its enigmatic ability to treat a wide range of illnesses, this valuable plant, which produced a gum-resin valued at an equivalent weight of silver, was well-known in Cyrene and sold to nearby nations.

Sylphium was certainly overexploited for many centuries, which resulted in its scarcity and eventual extinction by the fifth century A.D. [2]. In many societies, medicinal plants form the foundation of health care systems. Over 60% of the global population and 80% of those living in underdeveloped nations mostly rely on plants for their medicinal needs. The preservation of biodiversity, the development of new medications, and the improvement of the standard of living in impoverished rural communities are all tied to the recovery of knowledge and procedures related to these plant resources [2]. The therapeutic and medicinal qualities of different plants are utilized in herbal supplements, botanical-based nutraceuticals, and teas, in addition to serving as the foundation for between 30% and 40% of modern conventional medications [3].

Ethnobotanical research on medicinal plants has taken many different forms, sometimes detailing the usage of plants in specific cultural contexts and other times testing theories of use as well as knowledge [4]. Libya is one of the countries in Africa which have a variety of plants, especially in the eastern north side of Libya. Therefore, many studies were carried out to estimate many different compounds [5-38], besides the contents of minerals and metals were studied in different samples by using different methods [38-69]. This study aims to estimate some of the chemical constituents (Carbohydrates, total phenols, and antioxidants) in some selected plants. Using phytochemicals of leaves and stems. To measure the contents of the (minerals: Na, K, and Ca) in leaves and stems of *Capparis spinosa* L. (Kabbar), *Ceratonia siliqua* L. (Kharuwb), and *Juniperus phoenicea* L. (Arar) growing at Al-Gabal Al-Ahder regions, Libya.

Methods

Sampling

In this study, the leaves and stems of *Capparis spinosa* L. (Kabbar), *Ceratonia siliqua* L. (Kharuwb), and *Juniperus phoenicea* L. (Arar) plants were selected, where the samples were collected from different locations, including the valley called Wadi Derna, and Karsah in the West, Al-Dhahr Al-Ahmar in the South, and the Mediterranean coast in the North. The study region is situated on the second terrace of El-Jabal El-Akhdar Mountain, which is part of Wadi Derna in the Derna region of northeastern Libya. The Wadi separates the city into two halves between longitudes (33°00'–32° 30' N and 22°30'–22°45'E). The Wadi is between 40 and 300 meters above sea level. With a mean temperature of roughly 20°C, the research area's climate is similar to El-Jabal El-Akhdar. Between 200 and 300 mm of rain falls on average.

Sample extraction

10 grams of each dried sample were taken and transferred to a beaker containing 100 ml of distilled water, and the mixture was mixed. then the extraction was carried out by an evaporator system at 75 °C. The mixture was filtered. After two hours, the filtrate was used to determine the phytochemical screening [6-10].

Phytochemical Analysis

The central lab of the Faculty of Science at Omar Al Mukhtar University processed all of the phytochemical screening tests in accordance with the normal procedures, which are detailed in earlier research [11–15].

sterols and/or triterpines test: Libermann-Burchard's test

After adding 0.3 ml of acetic anhydride and 1 ml of the alcohol and aqueous extracts from each sample, only a few drops of concentrated sulphuric acid were applied through the side of dry test tubes. Once sterols and/or triterpines exist, a reddish-violet hue is created at the intersection of the two layers, and then the chloroform solution turns green.

Flavonoids test

The analyzed aqueous extracts, in addition to alcohol, were additionally extracted by using hydrochloric acid (1%). Each sample was then placed through the following test: 10ml from each extract was made alkaline, which produces a yellow when flavonoids are present.

Alkaloids test

The examined species' alcohol and aqueous extracts were extracted using 20 ml of diluted hydrochloric acid, kept cool, and made alkaline using a diluted ammonium hydroxide solution, followed by being extracted using chloroform.

The following tests are performed on the chloroform extract: Dragendorff's reagent preparation: Solution (a): A mixture of 10 ml acetic acid with 40 ml distilled aqueous was used to dissolve around 0.85 g of basic bismuth nitrate. Solution (b): 20 ml of water was used to dissolve about 8 grams of potassium iodide. Stock remedy: Solutions (a) and (b) are combined in equal volumes. Filter paper was treated with a few drops of chloroform extract, dried, and then sprayed with the reagent. When alkaloids are present, an orange hue is seen.

Tannins test

The solution of hydro-alcoholic clearis is put through the following test after the alcohol and aqueous extracts of the examined species were additionally extracted with 50% ethanol, followed by filtering out: Test for ferric chloride: The alcohol and aqueous solution were mixed with one ml of the reagent (1% FeCl₃). When pyrogallol tannins are present, a blue hue appears.

Anthraquinones test

Borntrück's test

After adding one ml each of the alcohol and aqueous extract of the subsequent extracts, either caustic soda or aqueous ammonia, the mixture is shaken. When anthraquinone glycosides are present, the aqueous layer takes on a rose-red hue.

Modified-Bornträger's test

Alcoholic potassium hydroxide is used to hydrolyse one ml each of the alcohol and aqueous extract of the subsequent extracts of the tested plants. This process is then acidified and continues as Borntrück's test. When anthraquinones are present, the aqueous layer turns rose-red.

Saponine test

After adding 5 ml of the tape aqueous solution to one ml of each alcohol and aqueous extract, the mixture is rapidly agitated for 5 minutes. The presence of saponin is indicated by the development of a foam that is one centimeter high and lasts for fifteen minutes.

Determination of Phenol Compounds by the Folin-Ciocalteu Method

The purpose of this experiment was to identify phenolic compounds. Using gallic acid as a reference, the content of total phenolics in the extracts was measured using the Folin-Ciocalteu reagent in accordance with the method of Slinkard and Singleton (10). Following the introduction of samples (two duplicates of each sample) into test cuvetts, 0.8 ml of Na₂CO₃ (7.5%) and 1.0 ml of Folin-Ciocalteu reagent were added. The Shimadzu UV-Vis spectrophotometer was used to measure the absorbance of each sample at 765 nm following a 1.5-hour incubation period at 30 °C. PPM of fresh weight was used to express the results.

Determination of antioxidant capacity by the Prussian blue method

One gram of the powdered substance was defatted using petroleum ether. The defatted powder has been extracted again using 10 ml of 1% hydrochloric acid: methanol (v/v) after two rounds of stirring with 10 ml of methanol. subsequently, those three combined extracts were vacuum-evaporated, and the residue was dissolved in 10 mL of methanol. Half a milliliter of the solution was mixed together with 3 distilled aqueous solutions, 3 ml of K₃Fe (CN)₆ (0.008 M), 3 ml of 0.1M HCl, and one milliliter of 1% FeCl₃.

The absorbance is measured at 720 nm in the central lab of Omar Al-Mukhtar University's Faculty of Science after the blue colour is allowed to develop for five minutes.

Determination of Carbohydrates

To estimate total carbohydrates, a known weight of 0.2 g of the dried sample was ground, then 5 mL of sulphuric acid was added. After completing, the samples were dissolved, the samples were cooled at room temperature, then a small quantity of Barium carbonate (Ba₂CO₃) was added, and the mixture was heated again. After cooling, the samples were filtered. One ml of solution was taken, then one ml of 5% phenol was added. The total carbohydrate was determined by the method carried out in a previous study. Where the absorbance was measured at a wavelength of 490 nm [8-10].

Determination of Minerals

In the Faculty of Science at Omar Al-Mukhtar University central lab, the contents of sodium, potassium, and calcium were measured using a Flame Photometer (JENWAY Flame Photometer) in accordance with the approach outlined by several researchers [60-70].

Results

This study verified the presence of sterols in all the selected samples. There is are relative increase in their contents in leaves compared with stems in the plants of *Capparis Spinosa*, *Ceratonia siliqua*, and *Juniperus phoenicea*. The high contents of flavonoids were recorded in leaves compared with the other samples of *Ceratonia siliqua* leaves and stems of the other plants. The results also recorded that the alkaloids were completely absent in *Capparis Spinosa* stems, while high content was detected in *Ceratonia siliqua* plant leaves and *Juniperus phoenicea* plant stems. Tannin compounds were recorded in all the studied samples except in *Capparis Spinosa* stems. Also, the Anthraquinones were not detected in the leaves and stems of the *Capparis Spinosa* plant samples. On the other side, the Anthraquinones showed relatively higher content in the leaves of the *Ceratonia siliqua* plant sample. The saponins were detected in all the selected samples of plants in this study. (Table 1&2).

Table 1. The phytochemical screening of sterols, flavonoids, and Alkaloids of the studied plants

Scientific name Compounds	Sterols		Flavonoids		Alkaloids	
	Leaves	Stems	Leaves	Stems	Leaves	Stems
<i>Capparis Spinosa</i> .	++	+	++	+	+	-
<i>Ceratonia siliqua</i> .	++	+	+++	+	+++	++
<i>Juniperus phoenicea</i>	++	++	+	++	+	+++

Table 2. The phytochemical screening of the studied plants

Scientific name Compounds	Tannins		Anthraquinones		Saponine	
	Leaves	Stems	Leaves	Stems	Leaves	Stems
<i>Capparis Spinosa</i> .	+	-	-	-	+++	+
<i>Ceratonia siliqua</i> .	+++	+++	+++	++	+++	+
<i>Juniperus phoenicea</i>	++	++	++	++	+++	++

Total phenols, Anti-Oxidant, and Carbohydrate Contents

The concentrations of total phenol contents showed variations between leaf and stem samples, fluctuating in the ranges of (222.72 -302.646 ppm) and (307.625 -319.013 ppm) of leaves and stems, respectively, (Table 3). A relative decrease in total phenol contents was observed in leaves compared with their values in *Ceratonia siliqua* and *Juniperus phoenicea* plants. The antioxidant activity, were ranged between (9.07 -10.59 ppm) and (9.955 -10.655 ppm), respectively. For the carbohydrate compounds, the results no recorded variations were observed between their contents in leaves compared with stems, where their values ranged between (0.039 -0.155 ppm) and (0.0016 – 0.029 ppm) were observed for anti-oxidant between stems and leaves.

Table 3. The contents (ppm) of Phenols, Anti-oxidant, and Carbohydrate in the studied samples

Scientific name Compounds	Total Phenols		Anti-Oxidant		Carbohydrate	
	Leaves	Stems	Leaves	Stems	Leaves	Stems
<i>Capparis Spinosa.</i>	302.646	319.0313	9.077	9.955	0.039	0.029
<i>Ceratonia siliqua.</i>	285.97	307.625	10.02	10.33	0.108	0.013
<i>Juniperus phoenicea</i>	222.72	318.549	10.595	10.655	0.155	0.0016

Minerals

The study results of this recorded presence of sodium, potassium and calcium in leaves and stems of the studied plants, the contents of sodium were ranged as following: (0.958 – 2.625 ppm) and (0.88 – 9.08 ppm) in leaves and stems, correspondingly, the results indicated that there as increase the sodium contents in *Capparis Spinosa* and *Ceratonia siliqua* plants comparing with *Juniperus phoenicea* plant, especially in stems, the concentrations of potassium showed higher levels comparing with sodium and calcium, also there are high concentrations in *Capparis Spinosa* plant comparing with *Ceratonia siliqua* and *Juniperus phoenicea* plants. The results of Minerals showed high contents of potassium in both leaves and stems of the studied samples, compared with the contents of Sodium and Calcium. (Table 4) The Potassium levels were ranged as (3.56 -77.16 ppm) for leaves and from (42.96- -76.96 ppm) for stems. For the calcium contents, the results of this study showed lower contents in leaves compared with stems of the studied plants, where the contents ranged between (0.08 -0.28 ppm) for leaves and between (0.375 -0.875 ppm), (Table 4).

Table 4. The contents (ppm) of minerals (Na, K, and Ca) in the studied samples

Scientific name Compounds	Sodium		Potassium		Calcium	
	Leaves	Stems	Leaves	Stems	Leaves	Stems
<i>Capparis Spinosa.</i>	2.625	9.08	77.16	76.96	0.28	0.541
<i>Ceratonia siliqua.</i>	0.958	4.08	6.16	48.16	0.08	0.375
<i>Juniperus phoenicea</i>	1.041	0.88	3.56	42.96	0.08	0.875

Discussion

The phytochemical screening in this study presented the presence of diverse amounts of natural compounds as flavonoids, alkaloids, sterols, tannins, phenols, Saponins, and anthraquinones, in the studied plants of *Ceratonia siliqua*, *Ceratonia siliqua*, and *Juniperus phoenicea*. The results showed variations in their contents between the studied plants, also between the leaves and stems. Colour tests were used to apply the deflection in accordance with conventional procedures. For the chemical analysis, most of the selected plants contain very important contents of Natural products, as sterols, phenols, Tannins, anthraquinones, and carbohydrates. Most of the selected plants have antioxidant and phenol compounds, which may be attributed to the presence of different aromatic chemical compounds.

Most medicinal plants contain benzene derivatives and different substitutions alkanes, which give special odors to most of them. Numerous secondary metabolites, including quinines, alkaloids, phenols, sterols, flavonoids, tannins, and saponins, are typically produced by plants and are significant sources of biocides and numerous other pharmaceutical medications. In pharmacological studies as well as drug development, plants with medicinal properties are crucial. The western pharmacopoeia contains about 7,000 therapeutic substances that originated from plants.

Given that they produce an extensive variety of bioactive chemicals, the majority of which are expected to originate as chemical defense against infection or predation and antioxidant substances, plants have been an excellent source of medications. Numerous plants have antibacterial properties, and these properties are employed to treat various illnesses [70]. Although they support an organism's proper growth and development, the organic substances created in the kingdom of plants do not directly contribute to the development, growth, or reproduction of organisms. Plant Ingredients that are biosynthetically produced from primary metabolites but have a more restricted distribution within the kingdom of plants are known

as secondary metabolites. carbohydrates, flavonoids, alkaloids, saponins, terpenes, tannins, lipids, and Phenolics are among the classes of secondary plant metabolites [71]. Therefore, open-access journals must motivate scientists and medical professionals to put in a lot of effort to identify the primary active components that can be isolated from plants for medicinal purposes. The variety and richness of electronic material about medicinal plants as a reemergent health aid has exploded due to recent and renewed interest in these plants, as well as advancements in information technology [72]. On the other side the main sources of minerals in the plants are coming from the areas around the plants as the agriculture soil and the climate or from the water [73-85], in this study high contents of potassium were observed comparing with sodium and calcium, generally most of natural plants containing higher potassium value due to its importance for the most plants. Also, there are chemical methods used to show the types the salts of minerals as XRD, XRF, ICP, and others. Some of these salts included minerals chlorides, minerals carbonate, minerals sulphate to explain their resources from sources a soils or coming from other sources as pollution of the sea water evaporator especially for the plants growth nearly of location near seas .some studies used different instrumental methods as XRF, atomic absorption, flame photometer to estimation the types and contents of minerals and metals in different natural and environmental samples included soil, sediment, water, plant, vegetable and others. Some studies have found some chemical compounds in natural samples as soils, plants, and waters, mainly coming from human activities [86-95]. Some studies used the metals in other applications as antibacterial agents and other uses [96-103].

Conclusion

The study's findings indicate that the chosen plants include a variety of natural compounds, including sterols, saponins, phenols, flavonoids, tannins, anthraquinones, and alkaloids, as well as differences in their contents in the leaves and stems. Small amounts of calcium were recorded compared with potassium and sodium. The results also showed that the studied plants contained different amounts of antioxidants and total phenols in both leaves and stems.

Acknowledgment

Grateful thanks to the staff of the chemistry department's central laboratory at Omar Al-Muhtar University's Faculty of Science for their assistance in setting up this study.

Conflict

The authors provide that no conflict in the results recorded in this, a contrast to other studies.

References

1. El-Mokasabi FM. The traditional use botanical composition of medicinal plants in the green mountain region [Doctoral dissertation]. [Alexandria, Egypt]: Alexandria University, Botany Department; 2010.
2. Almeida CF, Amorim EL, Albuquerque UP, Maia MB. Medicinal plants popularly used in the Xingo region – a semi-arid location in Northeastern Brazil. *J Ethnobiol Ethnomed*. 2006;2:15.
3. Chouaibi MN, Mahfoudhi L, Rezig F, Donsi G, Ferrari M, Hamdi S. Nutritional composition of *Zizyphus lotus* L. seeds. *J Sci Food Agric*. 2012;92(6):1171-7.
4. Borgi W, Chouchane N. Anti-spasmodic effects of *Zizyphus lotus* (L.) Desf. extracts on isolated rat duodenum. *J Ethnopharmacol*. 2009;126(3):571-3.
5. Abdull-Jalliel H, Sulayman A, Alhoreir M, Hasan H. Antimicrobial effect of some metal concentration on growth of staphylococcus and klebsiella bacteria species. *AlQalam J Med Appl Sci*. 2025;8(3):1646-56.
6. Abdull-Jalliel H, B Arous N, Alhoreir M, Hasan H. Using extracts of (Dodder) plant and concentrations of some metals as inhibitors for growth, (*Pseudomonas*) bacteria isolated from some hospital rooms in Derna and Al bayda. *AlQalam J Med Appl Sci*. 2025;8(3):1600-11.
7. Eltawaty SA, Abdalkader GA, Hasan HM, Houssein MA. Antibacterial activity and GC-MS analysis of chloroform extract of bark of Libyan *Salvia fruticosa* Mill. *Int J Multidiscip Sci Adv Technol*. 2021;1(1):715-21.
8. Aljamal MA, Hasan HM, Al Sonosy HA. Antibacterial activity investigation and anti-biotic sensitive's for different solvents (Ethanol, propanol, DMSO and di El er) extracts of seeds, leaves and stems of (*Laurus azorica* and *Avena sterilis*) plants. *Int J Curr Microbiol App Sci*. 2024;13(11):175-90.
9. Hamade MH, Abdelraziq SA, Gebreel AA. Extraction and determination of Beta carotene content in carrots and tomato samples collected from some markets at ElBeida City, Libya. *EPH Int J Appl Sci*. 2019;1(1):105-10.
10. Hasan HM, Ibrahim H, Gonaïd MA, Mojahidul I. Comparative phytochemical and antimicrobial investigation of some plants growing in Al Jabal Al-Akhdar. *J Nat Prod Plant Resour*. 2011;1(1):15-23.
11. Hasan H, Jadallah S, Zuhir A, Ali F, Saber M. Anti-Cancer, Anti-Inflammatory, Antibacterial, Antifungal, Anti-Oxidant and phytochemical investigation of flowers and stems of *Anacyclus Clavatus* plant extracts. *AlQalam J Med Appl Sci*. 2025;8(3):415-27.
12. Hasan H, Zuhir A, Shuib F, Abdraba D. Phytochemical investigation and exploring *Citrullus Colocynthis* extracts as antibacterial agents against some gram and negative bacteria species. *AlQalam J Med Appl Sci*. 2025;8(3):392-400.

13. MdZeyauallah R, Naseem A, Badrul I, Hamad MI, Azza SA, Faheem AB, et al. Catechol biodegradation by *Pseudomonas* strain: a critical analysis. *Int J Chem Sci.* 2009;7(3):2211-21.
14. El-Mehdawy MF, Eman KS, Hamad MI, Hasan H. Amino acids contents of leaves and stems for two types of herbal plants (Marjoram and Hybrid tea rose) at AL-Gabal AL-Akhder region. *Der Pharma Chem.* 2014;6(6):442-7.
15. El-Mehdawy MF, Eman KS, Hamad MIH. Amino acid contents of leaves and stems for three types of herbal plants at Al-Gabal Al-Akhder region. *World J Chem.* 2014;9(1):15-9.
16. Hamad MH, Noura AAM, Salem AM. Phytochemical screening, total phenolic, anti-oxidant, metal and mineral contents in some parts of plantago Albicans grown in Libya. *World J Pharm Res.* 2024;13(3):1-17.
17. Anees AS, Hamad MIH, Hasan H, Mojahidul I. Antifungal potential of 1,2-4triazole derivatives and rapeutic efficacy of *Tinea corporis* in albino rats. *Der Pharm Lett.* 2011;3(1):228-36.
18. Hamad H, Marwa M, Amal H. Determining contents of antioxidants, total phenols, carbohydrate, total protein, and some elements in *Eucalyptus gomphocephala* and *Ricinus communis* plant samples. *Libyan Med J.* 2015;1(1):222-31.
19. Hamad H, Zuhir A, Farag S, Dala A. Efficiency of *Cynara Cornigera* fruits on antibacterial, antifungal and its phytochemical, antioxidant screening. *Libyan Med J.* 2025;3(1):120-8.
20. Hamad H, Ashour S, Ahmed A. Estimation of amino acid composition, total carbohydrate, and total protein content in *Ballota pseudodictamnus* plant extracts from Al Jabal Al Akhdar Region, Libya. *Libyan Med J.* 2025;3(1):266-71.
21. Hamad H, Ahmed H, Wafa A. Evaluation of anti-oxidant capacity, total phenol, metal, and mineral contents of *Ziziphus lotus* plant grown at some regions of AlGabal AlKhder, Libya. *Libyan Med J.* 2025;3(1):137-43.
22. Ben Arous NA, Naser ME, Hamad MAH. Phytochemical screening, anti-bacterial and anti-fungal activities of leaves, stems and roots of *C. parviflorus* Lam and *C. salviifolius* L plants. *Int J Curr Microbiol App Sci.* 2014;13(11):262-80.
23. Anas FAE, Hamad MAH, Salim AM, Azza MH. Phytochemical screening, total phenolics, antioxidant activity and minerals composition of *Helichrysum stoechas* grown in Libya. *Afr J Biol Sci.* 2024;3(6):2349-60.
24. Naseer RE, Najat MAB, Salma AA, Hamad MAH. Evaluation of metal and mineral contents of leaves, stems and roots of *C. Parviflorus* Lam and *C. Salviifolius* L plants growing at Al Ghabal Al-Khder (Libya). *Int J Adv Multidiscip Res Stud.* 2024;4(5):191-4.
25. Hamad MAH, Salem AM. Total carbohydrate, total protein, minerals and amino acid contents in fruits, pulps and seeds of some cultivars of muskmelon and watermelon fruit samples collected from Algabab Alkhder region. *Sch J Appl Med Sci.* 2024;12(1):1-7.
26. Gonaïd MI, Ibrahim H, Al-Arefy HM. Comparative chemical and biological studies of *Salvia fruticosa*, *Ocimum basillicum* and *Pelargonium graveolans* cultivated in Al-Jabal Al- AkhdarAkhdar. *J Nat Prod Plant Resour.* 2012;6(2):705-10.
27. Rinya FMA, Hamad MAH, Ahlam KA, Hammida MEH. Phytochemical screening of some herbal plants (*Men*, *Origanum* and *Salvia*) growing at Al-Gabal Al-akhder Region-Libya. *Afr J Basic Appl Sci.* 2017;9(3):161-4.
28. Anas FAA, Hamad MAH, Salim AA, Azza MH. Phytochemical screening, total phenolics, antioxidant activity and minerals composition of *Helichrysum stoechas* grown in Libya. *Afr J Biol Sci.* 2024;3(6):2349-60.
29. Haroon A, Hasan H, Wafa AAS, Baset ESM. A comparative study of morphological, physiological and chemical properties of leaves and steam samples of (*E.gomphocephala*) (Tuart) plant growing at coastal (Derna city) and *J Res Environ Earth Sci.* 2024;9(12):10-8.
30. Hamad MAS, Ali AR. Separation and identification speciation of phenolic compounds in fruits and leaves of some medicinal plants (*Juniperus phoenicea* and *Quercus coccifera*) growing at Algabab Al Akhdar region, Libya. *Indian J Pharm Educ Res.* 2016;51(3):299-303.
31. Enam FM, Wesam FAM, Hamad MAH. Detection contents of minerals of (Sodium, Potassium and Calcium) and some metals of (Iron, Nickel and Copper) in some vegetable and the samples collected from Al-Marj. *Int J Adv Multidiscip Res Stud.* 2023;5(3):304-9.
32. Hamad MIH, Safa RM Mousa. synthesis and (IR and TEM) characterization of leaves and stem nanoparticles of *Artemisia* plant: comparative study for evaluation of anti-bacterial efficiency. *Int J Adv Multidiscip Res Stud.* 2024;4(5):195-9.
33. Elsalhin H, Abobaker HA, Hasan H, El-Dayek GA. Antioxidant capacity and total phenolic compounds of some algae species (*Anabaena* and *Spirulina platensis*). *Sch Acad J Biosci.* 2016;4(10):782-6.
34. Alaila AK, El Salhin HE, Ali RF, Hasan HM. Phytochemical screening of some herbal plants (*Men*, *Origanum* and *Salvia*) growing at al-gabal al-akhder region- Libya. *Int J Pharm Life Sci.* 2017;8(4):5500-3.
35. Hasan H, Mariea FFE, Eman KS. contents of some chemical compounds of leaves and stems of some herbal plants (*Thymy*, *Rosemary*, *Salvia*, *Marjoram* and *Hybrid Tea Rose*) at Al-Gabal Al-Akhder region. *EPH Int J Appl Sci.* 2014;6(3):1-8.
36. El-Mehdawe MF, Eman KS, Hamad MIH. Heavy metals and mineral elements contents of leaves and stems for some herbal plants at AL-Gabal AL-Akhder region. *Chem Sci Rev Lett.* 2014;3(12):980-6.
37. Hamad MIH, Aaza IY, Safaa SHN, Mabrouk MS. Biological study of transition metal complexes with adenine ligand. *Proc.* 2019;41(1):77.
38. Hasan JA, Hasan HMA. Potential human health risks assessment through determination of heavy metals contents in regularly consumed yogurta in Libya. *World J Pharm Pharm Sci.* 2024;13(12):100-12.
39. Mamdouh SM, Wagdi ME, Ahmed MA, Alaa EA, Essam AM, Hamad MIH. Rice husk and activated carbon for waste water treatment of El-Mex Bay, Alexandria Coast, Egypt. *Arab J Chem.* 2016;9(S2):S1590-S6.

40. Mamdouh SM, Wagdi ME, Ahmed MA, Alaa EA, Hamad IH. Heavy metals accumulation in sediments of Alexandria coastal areas. *Bull Fac Sci*. 2012;47(1-2):12-28.
41. Mamdouh SM, Wagdi ME, Ahmed MA, Hamad MIH. Chemical studies on Alexandria coast sediment. *Egypt Sci Mag*. 2005;2(4):93-102.
42. Mamdouh SM, Wagdi ME, Ahmed MA, Alaa EA, Hamad MIH. Distribution of different metals in coastal waters of Alexandria, Egypt. *Egypt Sci Mag*. 2010;7(1):1-19.
43. Mohamed AE, Afnan SA, Hamad MA, Mohammed AA, Mamdouh SM, Alaa RE, et al. Usage of natural wastes from animal and plant origins as adsorbents for removal of some toxic industrial dyes and heavy metals in aqueous media. *J Water Process Eng*. 2023;55:104192.
44. Mohamed HB, Mohammed AZ, Ahmed MD, Hamad MAH, Doaa AE. The heavy metal pollution and associated toxicity risk assessment in Ajdabiya and Zueitina, Libya. *Sci J Damietta Fac Sci*. 2024;14(1):16-27.
45. Nabil B, Hamad H, Ahmed E. Determination of Cu, Co and Pb in selected frozen fish tissues collected from Benghazi markets in Libya. *Chem Methodol*. 2018;2:56-63.
46. Wesam FAM, Hamad MAH. Detection of heavy metals and radioactivity in some bones of frozen chicken samples collected from Libyan markets. *Int J Adv Multidiscip Res Stud*. 2023;3(3):761-4.
47. Wesam FAM, Hamad MAH. Study the accumulation of minerals and heavy metals in *Ulva* algae, *Cladophora*, *Polysiphonia* and *Laurencia* algae samples at the eastern north region of the Libya coast. *GSC Biol Pharm Sci*. 2023;23(03):147-52.
48. Citrine E, Hamad H, Hajer Af. Contents of metal oxides in marine sediment and rock samples from eastern Libyan coast, utilizing X-ray method. *AlQalam J Med Appl Sci*. 2015;1(1):1316-21.
49. Hanan MA, Hamida E, Hamad MAH. Nitrogen, phosphorus and minerals (Sodium, Potassium and Calcium) contents of some algae's species (*Anabaena* and *Spirulina platensis*). *Int J Curr Microbiol App Sci*. 2016;5(11):836-41.
50. Mardhiyah F, Hamad H. Assessment of the contamination by heavy metals in Al-Fatayeh Region, Derna, Libya. *AlQalam J Med Appl Sci*. 2025;8(3):1081-91.
51. Abdelrazeg A, Khalifa A, Mohammed H, Miftah H, Hamad H. Using melon and watermelon peels for removal of some heavy metals from aqueous solutions. *AlQalam J Med Appl Sci*. 2025;8(3):787-96.
52. Abdul Razaq A, Hamad H. Estimate contents and types of water well salts by Palmer Roger model affecting corrosion of Al-Bayda city (Libya) network pipes. *AlQalam J Med Appl Sci*. 2025;8(3):744-53.
53. Abdulsayid FA, Hamad MAH, Huda AE. IR spectroscopic investigation, X-ray fluorescence scanning, and flame photometer analysis for sediments and rock samples of Al-Gabal Al-Akhder coast region (Libya). *IOSR J Appl Chem*. 2021;14(4):20-30.
54. ALambarki M, Hasan HMA. Assessment of the heavy metal contents in air samples collected from area extended between Albayda and Alquba cities (Libya). *AlQalam J Med Appl Sci*. 2025;8(3):695-707.
55. Al-Nayyan N, Mohammed B, Hamad H. Estimate of the concentrations of heavy metals in the and some plant samples collected from (near and far away) from the main road between Al-Bayda city and Wadi Al-Kouf region. *AlQalam J Med Appl Sci*. 2025;8(3):816-26.
56. Hasan HMI. Studies on physicochemical parameters and water treatment for some localities along the coast of Alexandria [Doctoral dissertation]. [Alexandria, Egypt]: Alexandria University; 2006.
57. Hamad MAH, Hager AA, Mohammed EY. Chemical studies of water samples collected from area extended between Ras Al-Halal and El Hania, Libya. *Asian J Appl Chem Res*. 2022;12(3):33-46.
58. Hamad M, Mohammed AA, Hamad MAH. Adsorption and kinetic study for removal some heavy metals by use in activated carbon from sea grasses. *Int J Adv Multidiscip Res Stud*. 2024;4(6):677-85.
59. Hamad MAH, Hamad NI, Mohammed MYA, Hajir OAA, Al-Hendawi RA. Using bottom marine sediments as environmental indicator state of the (Tolmaitha – Toukra) region at the eastern north coast of Libya. *Sch J Eng Tech*. 2024;2(14):118-32.
60. Hamad MIH. heavy metals distribution in the coastal water of Derna city (Libya). *Egypt J Aquat Res*. 2008;34(4):35-52.
61. Hamad MIH, Mojahid ul Islam. concentrations of some heavy metals in Al-Gabal Al-Akhder Coast Sediment. *Arch Appl Sci Res*. 2010;2(6):59-67.
62. Hamad MAH, Amira AKA. Estimate concentrations of some heavy metals in some shoes polish samples. *EPH Int J Appl Sci*. 2016;2(2):24-7.
63. Hamad MAH, Hussien SSM, Basit EEM. Accumulation of some heavy metals in green algae as bio indicators of environmental pollution at Al-Hania region: Libya coastline. *Int J Adv Multidiscip Res Stud*. 2024;4(5):188-90.
64. Hamad MIH, Ahmed MA. Major cations levels studies in surface coastal waters of Derna city, Libya. *Egypt J Aquat Res*. 2009;35(1):13-20.
65. Hamad MIH, Masoud MS. rmal analysis (TGA), diffraction rmal analysis (DTA), infrared and X-rays analysis for sediment samples of Toubrouk city (Libya) coast. *Int J Chem Sci*. 2014;12(1):11-22.
66. Hamad R, Ikraiam FA, Hasan H. Estimation of heavy metals in the bones of selected commercial fish from the eastern Libyan coast. *J Rad Nucl Appl*. 2024;9(1):47-51.
67. Hasan HAH. Estimate lead and cadmium contents of some archeological samples collected from ancient cities location (Cyrene and Abolonia) at Al-Gabal Al-Akhder Region, Libya. *Univ J Chem Appl*. 2021;12(21):902-7.
68. Hasan H, El-maleh C. Evaluation of some heavy metal levels in tissue of fish collected from coasts of susa region, libya. *Attahadi Med J*. 2025;1(1):118-22.
69. Balal A, Obid M, Khatab H, Hasan H. Determination of lead and cadmium marine water and crabs (*pachygrapsus marmoratus*) from tolmitha coast, Libya. *AlQalam J Med Appl Sci*. 2025;8(3):1670-7.
70. Arora D, Kaur J. Antimicrobial activity of spices. *Int J Antimicrob Agents*. 1999;12(3):257-62.

71. Bhat KS. Medicinal and plant information databases. In: Medicinal Plants for Forests Conservation and Health Care. Rome: FAO; 2011. p. 158.
72. Anulika NP, Ignatius EO, Aymond ES, Osasere OL, Abiola AH. The chemistry of natural product; Plant secondary metabolites. *Int J Technol Eng Res.* 2016;4(8):1-9.
73. Hamad IH, Nuesry MS. poly cyclic hydrocarbons levels in some fishes tissues collected from Derna City (Libya) Coast. In: International conference on chemical, agricultural and medical sciences; 2014 Dec 4-5; Antalya, Turkey. 2014. p. 52-6.
74. Hamad MAH, Mounera AAE, Baseet ESM, Eman E, Al-Badri M. Identification and detection aromatic and aliphatic hydrocarbons in *Epinephelus Marginatus* fish samples collected from Benghazi coast. *Int J Adv Multidiscip Res Stud.* 2023;6(3):107-13.
75. Mohammed A, Hamad MAH, Mounera AAE, Eman IHE. Extraction and identification of aliphatic hydrocarbons in marine sediment samples at Benghazi city and Dyriana town coasts (Libya). *J Res Humanit Soc Sci.* 2023;11(10):168-74.
76. Hasan MAH, Muftah HS, Abdelghani KA, Saad SI. Poly aromatic hydrocarbon concentrations in some shell samples at some Tobrouk city coast regions: could the oil industry be significantly affecting the environment. *Ukr J Ecol.* 2022;12(3):21-8.
77. Habel AMA, Mohamed NIH, Mohammed MA, Hamad MAH. The levels and sources of aliphatic and polycyclic aromatic hydrocarbons in blue runner fish from Benghazi coast, Libya. *Afr J Biol Sci.* 2024;6(3):1-10.
78. Hasan HMI, Mohamad ASA. A study of aliphatic hydrocarbons levels of some waters and sediments at Al-Gabal Al-Akhder coast regions. *Int J Chem Sci.* 2013;11(2):833-49.
79. Salem GM, Aljidaemi FF, Hwisa SA, Hamad MIH, Zaid AA, Amer IO. Occupational exposure to benzene and changes in hematological parameters in East Tripoli, Libya. *Nanotechnol Percept.* 2024;20(S5):358-64.
80. Habil Z, Ben arous N, Masoud N, Hasan H. Using GC-mass method for determination hydrocarbon compounds in some vegetable samples at Derna city, Libya. *Libyan Med J.* 2025;17(3):374-83.
81. Hasan H, Habil Z, Ben arous N. Estimate the types and contents of phenolic acid in *C.Paviflorus lam* and *C.salviifolius L* plants growing at Al -Gabal Al-hder regions. *AlQalam J Med Appl Sci.* 2025;8(3):1646-56.
82. Zeyadah MA, Bahnasaway MH, Deedah AM, El-Emam DA, Hamad MA Hasan. Evaluation of the contents of aliphatic and aromatic hydrocarbons in sediment from Zwwitina harbor coast (Libya), ad indicator of petroleum pollution. *Egypt J Aquat Biol Fish.* 2023;27(6):1-18.
83. Hamad R, Ikraiam F, Hasan H. Determination of specific natural radionuclides in the bones of some local fish commonly consumed from the eastern Libyan coast. *J Rad Nucl Appl.* 2023;8(3):283-9.
84. Sroor AT, Walley El-Dine N, El-Bahi SM, Hasa HMA, Ali JM. Determination of radionuclides levels and absorbed dose for the, rock, plant and water in gondola- Libya. *IOSR J Appl Phys.* 2018;10(4):40-9.
85. Hasan H, Ammhmmid R, Khatib H, Ali J, Al kaseh A. Using gamma ray radiation to estimate the types and contents of radioactive nuclides in some ported sugar samples, Libya. *AlQalam J Med Appl Sci.* 2025;8(3):1795-803.
86. Hasan H, Abdelgader I, Emrayed H, Abdel-Gany K. Removal of the medical dye safranin from aqueous solutions by sea grasses activated carbon: a kinetic study. *AlQalam J Med Appl Sci.* 2025;8(3):428-34.
87. Hasan HMA, Alhamdy MA. Adsorption and kinetic study for removal some heavy metals by using activated carbon of sea grasses. *Int J Adv Multidiscip Res Stud.* 2024;4(6):677-85.
88. Almadani EA, Hamad MAH, Kwakab FS. Kinetic study of the adsorption of the removal of bromo cresol purple from aqueous solutions. *Int J Res Granthaalayah.* 2019;7(12):1-10.
89. Mamdouh SM, Wagdi ME, Ahmed MA, Alaa EA, Essam AM, Hamad MIH. Rice husk and activated carbon for waste water treatment of El-Mex Bay, Alexandria Coast, Egypt. *Arab J Chem.* 2016;9(S2):S1590-S6.
90. Alfutisi H, Hasan H. Removing of thymol blue from aqueous solutions by pomegranate peel. *EPH Int J Appl Sci.* 2019;1(1):111-9.
91. Ahmed ONH, Hamad MAH, Fatin ME. Chemical and biological study of some transition metal complexes with guanine as ligand. *Int J New Chem.* 2023;10(3):172-83.
92. Hamad MAH, Enas UE, Hanan AK, Hana FS, Somia MAE. Synsis, characterization and antibacterial applications of compounds produced by reaction between Barbitol with Threonine, glycine, lycine, and alanine. *Afr J Biol Sci.* 2024;6(4):1-10.
93. Emrayed H, Hasan H, Liser R. Corrosion inhibition of carbon steel using (Arginine -levofloacin-metal) complexes in acidic media. *AlQalam J Med Appl Sci.* 2025;8(3):1633-40.
94. Hasan S, Abduljalil O, Mohamed F, Hasan H. Detection of residual pesticides (Imidacloprid ,Aldicarb,Metalaxyl,Cypermethrin ,Chlorpyrfos,DDA, and Endrin) in peach Samples collected from Jabal al Akhder farma,Libya. *AlQalam J Med Appl Sci.* 2025;8(4):2099-106.
95. Mohamed FH, Salah MIH, Omuthum A, Hasan Hamad. Sensitive and rabid method to estimate residual pesticides in some local and imported apple cultivars collected from eastern north side of Libya. *Int J Adv Multidiscip.* 2023;3(6):100.
96. Hamad MIH, Aaza IY, Safaa SH, Mabrouk MS. Biological Study of Transition Metal Complexes with Adenine Ligand. *Proceedings.* 2019;41(1):77.
97. Ibrahim HH, Ahlam MAI, Hamad MIH. Synthesis, Infrared (I.R), Thermal Gravimetric Analysis (TGA) Characterization and Antibacterial Activity of some Amino Acids Complexes. *Chem Sci Rev Lett.* 2014;3(12):1303-16.
98. Hamad MAH, Hanan AAK, Fatima A. Infrared (IR) charactizations and physicochemical properties of Schiff base compound obtained by the reaction between 4-hydroxy-3-methoxy benzaldehyde and 2-amino-3-methylbutanoic acid. *J Res Pharm Sci.* 2021;7:8-12.

99. Mabrouk MS, Moussa SF, Hasan HA. Preparation, Characterization and Antibacterial Activities of Metal complexes with Tyrosine Ligands. Egypt J Chem. 2023;66(10):1-7.
100. Zawia A, Adel MN, Ashraf AAA, Tahani A. synthesis, Characterization, Biological Screening and Molecular Docking of New Schiff Base and Its Mononuclear Complexes with Pb²⁺, Cd²⁺, Zn²⁺ and Cu²⁺. J Pharm Appl Chem. 2023;8(1):1-9.
101. Hamad MAH, Fatin E, Ahmed O, Ahmed NH. Chemical and Biological Study of Some Transition Metal Complexes with Guanine as Ligand. Int J New Chem. 2023;10(3):172-83.
102. Hamad MAH, Yassmeen A. Synthesis, physical properties, Infrared (IR) analysis and anti-fungi activity of some Valine and metal ion Complexes. Int J Multidiscip Sci Adv Technol. 2021;1:246-54.
103. Hanan MMA, Hamad MIH. Removing of Thymol Blue from aqueous solutions by Pomegranate peel. EPH Int J Appl Sci. 2019;1(1):111-9.